

# Australian Bulk Alliance

A.B.N 39 087 280 260

## CANOLA RECEIVAL STANDARDS – 2001/2002 SEASON REFERENCE CHART



Grade	Moisture	Test weight	Foreign material By weight	Wild turnip By count	Broken seeds By count	Total defective		Damaged seeds (bin burnt, frosted, weather damaged) By weight	Sprouted seeds By count	Green seeds By count
TEST CODE	MO*	TW	TF	WT	BK	DG		DS	SP	GN
CANO	Max 8.0%	Min 62	Max 3.0%	Max 3.0%	Max 7.0%	Max 10.0%	of	Max 3.0%	Max 5.0%	Max 2.0%
CAN 2	Max 8.0%	Min 62	Max 3.0%	Max 3.0%	Max 7.0%	Max 20.0%	which	Max 5.0%	Max 5.0%	Max 10.0%

\*Total Defective is the addition of Damaged Seeds, Sprouted Seeds, Green Seeds

**There is a limit of no more than 1 snails/stones or equivalent in a ½ litre passing through a 3.00mm round hole screen.  
There is a NIL tolerance for any snails/stones or equivalent remaining above a 3.00mm screen (2.25 litre sample size).**

### Restricted Seed Contaminants

(Limit per 1/2 litre Sample )

Burrs (Xanthium Spp)	1	Jute	3
Common Heliotrope	3	Thornapple	2
Crow Garlic	2	Nightshade	3
Sesbania Pea	60		
Skeleton Weed	2		
Hexham Scent	3		
Mexican Poppy	3		
Darnel	3		
Mint Weed	3		

### Nil Tolerance Seed Contaminants

Any other Tainting Agents, or contaminant imparting an objectionable odour.

Alligator Weed	<i>Alteranthera philoxeroides</i>	Rattlepod	<i>Crotalaria spp</i>
Cape Tulip	<i>Homeria spp</i>	Saffron Thistle	<i>Carthamus lanatus</i>
Castor Oil Plant	<i>Ricinus communis L</i>	Dodder	<i>Cuscuta spp</i>
Creeping Knapweed	<i>Acoptilon repens</i>	Giant Sensitive Plant	<i>Mimosa invisia</i>
Darling Pea	<i>Swansonia spp</i>	Star Burr	<i>Acanthospermum hispidum</i>
Parthenium Weed	<i>Parthenium hysterophorus</i>	Stinkwort	<i>Inula graveolens</i>
Ragweed	<i>Ambrosia spp</i>	St Johns Wort	<i>Hypericum perforatum L</i>
Opium Poppy	<i>Papaver somniferum L</i>	Wild Mignonette	<i>Reseda lutea</i>
Coriander	<i>Coriandrum sativum</i>		